## 包裝異動公告-Augmentin Syrup 457mg/5ml 安滅菌糖漿用粉劑 457 毫克/5 毫升 新包裝圖示 項目 Im 8\gm 78\$ nitnamguA WAN WAN STATE GGK GSK Augmentin Before reconstitution, keep tightly closed and store in a dry place at 30°C or below. Augmentin When reconstituted as directed, each 5 ml contains 400 mg amoxicilin as the tilhydrate and 57mg clavulanic acid as the potassium salt. amoxicilin + davulanic acid Once reconstituted, the suspension must be stored in a refrigerator and used within 7 days. Contains aspartame: 糖學用粉幣457毫克/5毫升 For oral use only. 外 Twice Daily Dosing Do not freeze 工業員の製計費の役割 南藤電航年第12247世 銀幣 日本電車車工業業を有限公司 出土・セス・イン・ には、12年上旬間一般に対し場 開連車 Gao Wiktone Poliuston 展生、2 cs 1 Proteines, 53100 Mayerne, hance To be used as directed by your physician. 457 mg/5 ml Keep out of the reach of children. Manufactured by: Glaxo Wellcome Production 35 ml 配製方法 - 天成句称末 - 東/冷開水框句 - 再加速量(末里 - 東/冷開水框句 - 最後出海冷開水里根上記號( - ) - 最初性便得。 mixed fruit flavoured Trade marks are owned by or licensed to the GSK group of companies. 62022 GSK group of companies or its. syrup in powder form 4、改重冰板冷藏(2-8°C)復有、色調 在7日内月史+ 2 住存期間数色音信情数応信息只 要犯7日期間内皆可使用 ◆等次回用前間被逐進句。 ◆管色機構提句再使用。 Augmentin amoxicillin + clavulanic acid 6-3000000000859-18 Twice Daily Dosing 457 mg/6 ml mised truit flavoured / syrup in powder form



臺北市立聯合醫院藥劑部 依據中華民國 113 年 1 月 4 日廠商來函公告,包裝更新時間點依廠商配貨及各院使用情形而異。(112-015)