



藥物成分	劑型	櫃號
Rosuvastatin	口服藥	

☞ 藥品商品名及含量

- 諾脂替膜衣錠 10 毫克/錠 (Roty F.C.)
- 洛脂平膜衣錠 10 毫克/錠 (Rotlip F.C.)
- 樂脈舒膜衣錠 10 毫克(ROSU F.C. TABLETS 10MG)
- 柔舒脂膜衣錠 10 毫克(ROSULIP F.C. TABLETS 10MG)
- 冠脂妥膜衣錠 20 毫克(CRESTOR 20MG FILM-COATED TABLETS)

☞ 藥品作用

本藥可減慢體內膽固醇的生成，主要用來治療高膽固醇血症及高三酸甘油酯血症，減少發生心血管疾病或中風的機會。

☞ 服用方式

1. 通常每日 1 次，每次 0.5~1 粒。
2. 建議與制酸劑(胃藥，成分多為鋁、鎂製劑)隔開 2 個小時使用，避免影響本藥吸收。
3. 請依照醫師指示服藥，不可自行調整劑量或增加服藥次數。若不慎忘記服藥，請立即補服，但若已接近下一次服藥時間，在下次用藥時間服藥即可，不可為了補吃而服用雙倍劑量。

☞ 注意事項

1. 請告訴醫師或藥師您是否有藥物過敏或是有其他疾病，特別是**肝臟或腎臟**方面的疾病。
2. 請告訴醫師或藥師您是否有在服用其他藥物(包含健康食品)，特別是抗凝血劑 warfarin、cyclosporine、erythromycin 或其他降血脂藥。
3. 請勿飲酒及大量葡萄柚汁，可能會加重此藥之副作用。
4. 本藥可能會影響血糖控制，使用期間請注意血糖的變化。
5. 副作用相當少見，但有可能發生：**頭痛、消化不良、噁心、腹痛**，但一般都不嚴重，若症狀惡化且無法忍受請回診。
6. 如果有下列症狀，請立刻就醫：**若出現眼睛/皮膚變黃、**

臺北市立

聯合醫院

TAIPEI CITY

HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

TAIPEI CITY HOSPITAL

茶色尿液(黃疸)、右上腹疼痛、不明原因的肌肉疼痛、肌肉無力或痙攣，尤其伴有全身不適、虛弱或發燒的現象。。

7. 懷孕等級 X，女性計畫懷孕或已懷孕請先告知醫師；服藥期間不可哺餵母乳，有需要哺乳請事先諮詢醫師/藥師。